

Applications fractional calculus special operators

Ali Omar

1. INTRODUCTION, DEFINITIONS, AND PRELIMINARIES

Let \mathcal{S} denote the class of (normalized) functions of the form:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

which are analytic and univalent in the open unit disc $\mathcal{U} = \{z: z \in \mathbb{C} \text{ and } |z| < 1\}$. A function $f(z)$ in \mathcal{S} is said to be starlike of order α in \mathcal{U} if and only if

$$\Re \left\{ \frac{zf'(z)}{f(z)} \right\} > \alpha, \quad (z \in \mathcal{U}; 0 \leq \alpha < 1).$$

We denote by $\mathcal{S}^*(\alpha)$ the class of all functions in \mathcal{S} which are starlike of order α in \mathcal{U} .

A function $f(z)$ in \mathcal{S} is said to be convex of order α in \mathcal{U} if and only if

$$\Re \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \alpha, \quad (z \in \mathcal{U}; 0 \leq \alpha < 1).$$

The present investigation was supported, in part, by the Natural Sciences and Engineering Research Council of Canada under Grant OGP0007353. And we denote by $\mathcal{K}(\alpha)$ the class of all functions in \mathcal{S} which are convex of order α in \mathcal{U} . We note that

$$f(z) \in \mathcal{K}(\alpha) \Leftrightarrow zf'(z) \in \mathcal{S}^*(\alpha), \quad (0 \leq \alpha < 1).$$

The classes $\mathcal{S}^*(\alpha)$ and $\mathcal{K}(\alpha)$ were first introduced by Robertson [1], and were studied subsequently by Schild [2], MacGregor [3], and Pinchuk [4] (see also [5]).

The function

$$S_\alpha(z) = \frac{z}{(1-z)^{2(1-\alpha)}}, \quad (z \in \mathcal{U}; 0 \leq \alpha < 1)$$

is the well-known extremal function for $\mathcal{S}^*(\alpha)$. Setting

$$c_n(\alpha) = \frac{\prod_{k=2}^n (k - 2\alpha)}{(n-1)!}, \quad (n \in \mathbb{N} \setminus \{1\}; \mathbb{N} = \{1, 2, 3, \dots\})$$

$S_\alpha(z)$ can be written in the form:

$$S_\alpha(z) = z + \sum_{n=2}^{\infty} c_n(\alpha) z^n.$$

Then we can see that $c_n(\alpha)$ is a decreasing function in α ($0 \leq \alpha < 1$) and that

$$\lim_{n \rightarrow \infty} c_n(\alpha) = \begin{cases} \infty, & \left(\alpha < \frac{1}{2}\right), \\ 1, & \left(\alpha = \frac{1}{2}\right), \\ 0, & \left(\alpha > \frac{1}{2}\right). \end{cases}$$

Let $f * g(z)$ denote the Hadamard product of two functions $f(z)$ and $g(z)$, that is, if $f(z)$ is given by (1.1) and $g(z)$ is given by

$$g(z) = z + \sum_{n=2}^{\infty} b_n z^n$$

then

$$f * g(z) = z + \sum_{n=2}^{\infty} a_n b_n z^n.$$

Let $\mathcal{R}(\alpha, \beta)$ be the subclass of \mathcal{S} consisting of functions $f(z)$ such that

$$f * \mathcal{S}_\alpha(z) \in \mathcal{S}^*(\beta), \quad (0 \leq \alpha < 1; 0 \leq \beta < 1)$$

Further, let $\mathcal{C}(\alpha, \beta)$ be the subclass of \mathcal{S} consisting of functions $f(z)$ such that

$$zf'(z) \in \mathcal{R}(\alpha, \beta), \quad (0 \leq \alpha < 1; 0 \leq \beta < 1).$$

The class $\mathcal{R}(\alpha, \beta)$ of α -prestarlike functions of order β was introduced by Sheil-Small et al. [6].

Let \mathcal{T} denote the subclass of \mathcal{S} consisting of functions of the form:

$$f(z) = z - \sum_{n=2}^{\infty} a_n z^n, \quad (a_n \geq 0).$$

We denote by $\mathcal{R}[\alpha, \beta]$ and $\mathcal{C}[\alpha, \beta]$ the classes obtained by taking intersections, respectively, of the classes $\mathcal{R}(\alpha, \beta)$ and $\mathcal{C}(\alpha, \beta)$ with the class \mathcal{T} ; that is,

$$\mathcal{R}[\alpha, \beta] := \mathcal{R}(\alpha, \beta) \cap \mathcal{T}$$

and

$$\mathcal{C}[\alpha, \beta] := \mathcal{C}(\alpha, \beta) \cap \mathcal{T}$$

The class $\mathcal{R}[\alpha, \beta]$ was studied recently by Silverman and Silvia [7] and Uralegaddi and Sarangi [8]. The class $\mathcal{C}[\alpha, \beta]$ was studied by Owa and Uralegaddi [9].

The following known results will be required in our investigation of the classes $\mathcal{R}[\alpha, \beta]$ and $\mathcal{C}[\alpha, \beta]$

LEMMA 1. (Silverman and Silvia [7]). Let the function $f(z)$ be defined by (1.11). Then $f(z)$ is in the class $\mathcal{R}[\alpha, \beta]$ if and only if

$$\sum_{n=2}^{\infty} (n - \beta) c_n(\alpha) a_n \leq 1 - \beta.$$

The result is sharp.

LEMMA 2. (Owa and Uralegaddi [9]). Let the function $f(z)$ be defined by (1.11). Then $f(z)$ is in the class $\mathcal{C}[\alpha, \beta]$ if and only if

$$\sum_{n=2}^{\infty} n(n - \beta) c_n(\alpha) a_n \leq 1 - \beta$$

The result is sharp.

2. OPERATORS OF FRACTIONAL CALCULUS

We begin by recalling the following definitions of operators of fractional calculus (that is, fractional derivatives and fractional integrals) which were used by Owa [10,11] and (more recently) by Srivastava and Owa [12] (see also [13, p. 343]).

Definition 1. The fractional integral of order λ is defined, for a function $f(z)$, by

$$D_z^{-\lambda} f(z) = \frac{1}{\Gamma(\lambda)} \int_0^z \frac{f(\zeta)}{(z - \zeta)^{1-\lambda}} d\zeta, \quad (\lambda > 0),$$

where $f(z)$ is an analytic function in a simply-connected region of the z -plane containing the origin, and the multiplicity of $(z - \zeta)^{\lambda-1}$ is removed by requiring $\log(z - \zeta)$ to be real when $z - \zeta > 0$.

DEFINITION 2. The fractional derivative of order λ is defined, for a function $f(z)$, by

$$D_z^{\lambda} f(z) = \frac{1}{\Gamma(1 - \lambda)} \frac{d}{dz} \int_0^z \frac{f(\zeta)}{(z - \zeta)^{\lambda}} d\zeta, \quad (0 \leq \lambda < 1),$$

where $f(z)$ is constrained, and the multiplicity of $(z - \zeta)^{-\lambda}$ is removed, as in Definition 1.

Definition 3. Under the hypotheses of Definition 2, the fractional derivative of order $n + \lambda$ is defined by

$$D_z^{n+\lambda} f(z) = \frac{d^n}{dz^n} D_z^{\lambda} f(z), \quad (0 \leq \lambda < 1; n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}).$$

Our first set of distortion properties of the class $\mathcal{R}[\alpha, \beta]$, involving the fractional integral operator $D_z^{-\lambda}$, is contained in the following theorem. THEOREM 1. Let the function $f(z)$ defined by (1.11) be in the class $\mathcal{R}[\alpha, \beta]$ with $0 \leq \alpha \leq 1/2$ and $0 \leq \beta < 1$. Then

$$|D_z^{-\lambda} f(z)| \geq \frac{|z|^{1+\lambda}}{\Gamma(2+\lambda)} \left\{ 1 - \frac{1-\beta}{(2-\beta)(1-\alpha)(2+\lambda)} |z| \right\}$$

and

$$|D_z^{-\lambda} f(z)| \leq \frac{|z|^{1+\lambda}}{\Gamma(2+\lambda)} \left\{ 1 + \frac{1-\beta}{(2-\beta)(1-\alpha)(2+\lambda)} |z| \right\}$$

for $\lambda > 0$ and $z \in \mathcal{U}$. The results are sharp.

Proof. Let

$$\begin{aligned} F(z) &= \Gamma(2+\lambda) z^{-\lambda} D_z^{-\lambda} f(z) \\ &= z - \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(2+\lambda)}{\Gamma(n+1+\lambda)} a_n z^n \\ &= z - \sum_{n=2}^{\infty} \Phi(n) a_n z^n, \end{aligned}$$

where

$$\Phi(n) = \frac{\Gamma(n+1)\Gamma(2+\lambda)}{\Gamma(n+1+\lambda)}, \quad (\lambda > 0; n \in \mathbb{N} \setminus \{1\})$$

We note that

$$0 < \Phi(n) \leq \Phi(2) = \frac{2}{2+\lambda}, \quad (\lambda > 0; n \in \mathbb{N} \setminus \{1\})$$

and that

$$c_{n+1}(\alpha) \geq c_n(\alpha), \quad \left(0 \leq \alpha \leq \frac{1}{2}; n \in \mathbb{N} \setminus \{1\} \right)$$

by means of (1.6). Consequently, by using Lemma 1, we have

$$(2-\beta)c_2(\alpha) \sum_{n=2}^{\infty} a_n \leq \sum_{n=2}^{\infty} (n-\beta)c_n(\alpha) a_n \leq 1-\beta$$

which implies that

$$\sum_{n=2}^{\infty} a_n \leq \frac{1-\beta}{2(2-\beta)(1-\alpha)}$$

Therefore, by using (2.8) and (2.9), we find that

$$\begin{aligned}
|F(z)| &\geq |z| - \Phi(2)|z|^2 \sum_{n=2}^{\infty} a_n \\
&\geq |z| - \frac{1-\beta}{(2-\beta)(1-\alpha)(2+\lambda)} |z|^2
\end{aligned}$$

and

$$\begin{aligned}
|F(z)| &\leq |z| + \Phi(2)|z|^2 \sum_{n=2}^{\infty} a_n \\
&\leq |z| + \frac{1-\beta}{(2-\beta)(1-\alpha)(2+\lambda)} |z|^2,
\end{aligned}$$

which prove the inequalities (2.4) and (2.5) of Theorem 1.

The equalities in (2.4) and (2.5) are attained by the function $f(z)$ given by

$$D_z^{-\lambda} f(z) = \frac{z^{1+\lambda}}{\Gamma(2+\lambda)} \left\{ 1 - \frac{1-\beta}{(2-\beta)(1-\alpha)(2+\lambda)} z \right\}$$

or, equivalently, by

$$f(z) = z - \frac{1-\beta}{2(2-\beta)(1-\alpha)} z^2.$$

Thus, we complete the proof of Theorem 1.

In the same way, we can prove the following distortion theorem for the class $\mathcal{C}[\alpha, \beta]$ by using Lemma 2 instead of Lemma 1.

THEOREM 2. Let the function $f(z)$ defined by (1.11) be in the class $\mathcal{C}[\alpha, \beta]$ with $0 \leq \alpha \leq 1/2$ and $0 \leq \beta < 1$. Then

$$|D_z^{-\lambda} f(z)| \geq \frac{|z|^{1+\lambda}}{\Gamma(2+\lambda)} \left\{ 1 - \frac{1-\beta}{2(2-\beta)(1-\alpha)(2+\lambda)} |z| \right\}$$

and

$$|D_z^{-\lambda} f(z)| \leq \frac{|z|^{1+\lambda}}{\Gamma(2+\lambda)} \left\{ 1 + \frac{1-\beta}{2(2-\beta)(1-\alpha)(2+\lambda)} |z| \right\}$$

for $\lambda > 0$ and $z \in \mathcal{U}$. The results are sharp for the function $f(z)$ given by

$$D_z^{-\lambda} f(z) = \frac{z^{1+\lambda}}{\Gamma(2+\lambda)} \left\{ 1 - \frac{1-\beta}{2(2-\beta)(1-\alpha)(2+\lambda)} z \right\}$$

or, equivalently, by

$$f(z) = z - \frac{1 - \beta}{4(2 - \beta)(1 - \alpha)} z^2.$$

Next, we state and prove the following theorem.

THEOREM 3. Let the function $f(z)$ defined by (1.11) be in the class $\mathcal{R}[\alpha, \beta]$ with $0 \leq \alpha \leq 1/2$ and $0 \leq \beta < 1$. Then

$$|D_z^\lambda f(z)| \geq \frac{|z|^{1-\lambda}}{\Gamma(2-\lambda)} \left\{ 1 - \frac{1-\beta}{(2-\beta)(1-\alpha)(2-\lambda)} |z| \right\}$$

and

$$|D_z^\lambda f(z)| \leq \frac{|z|^{1-\lambda}}{\Gamma(2-\lambda)} \left\{ 1 + \frac{1-\beta}{(2-\beta)(1-\alpha)(2-\lambda)} |z| \right\}$$

for $0 \leq \lambda < 1$ and $z \in U$. The results are sharp.

Proof. Let

$$\begin{aligned} G(z) &= \Gamma(2-\lambda) z^\lambda D_z^\lambda f(z) \\ &= z - \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(2-\lambda)}{\Gamma(n+1-\lambda)} a_n z^n \\ &= z - \sum_{n=2}^{\infty} \Psi(n) n a_n z^n \end{aligned}$$

where

$$\Psi(n) = \frac{\Gamma(n)\Gamma(2-\lambda)}{\Gamma(n+1-\lambda)}, \quad (0 \leq \lambda < 1; n \in \mathbb{N} \setminus \{1\}).$$

From (2.9) and Lemma 1, it follows that

$$2(1-\alpha) \sum_{n=2}^{\infty} n a_n \leq (1-\beta) + 2\beta(1-\alpha) \sum_{n=2}^{\infty} a_n \leq \frac{2(1-\beta)}{2-\beta}.$$

Consequently, we have

$$\sum_{n=2}^{\infty} n a_n \leq \frac{1-\beta}{(2-\beta)(1-\alpha)}$$

Note also that

$$0 < \Psi(n) \leq \Psi(2) = \frac{1}{2-\lambda}, \quad (0 \leq \lambda < 1; n \in \mathbb{N} \setminus \{1\})$$

Therefore, by using (2.22) and (2.23), we observe that

$$\begin{aligned}
|G(z)| &\geq |z| - \Psi(2)|z|^2 \sum_{n=2}^{\infty} na_n \\
&\geq |z| - \frac{1-\beta}{(2-\beta)(1-\alpha)(2-\lambda)} |z|^2
\end{aligned}$$

and

$$\begin{aligned}
|G(z)| &\leq |z| + \Psi(2)|z|^2 \sum_{n=2}^{\infty} na_n \\
&\leq |z| + \frac{1-\beta}{(2-\beta)(1-\alpha)(2-\lambda)} |z|^2.
\end{aligned}$$

Thus, the proof of Theorem 3 is completed upon noting that the equalities in (2.18) and (2.19) are attained by the function $f(z)$ defined by

$$D_z^\lambda f(z) = \frac{z^{1-\lambda}}{\Gamma(2-\lambda)} \left\{ 1 - \frac{1-\beta}{(2-\beta)(1-\alpha)(2-\lambda)} z \right\}$$

or, equivalently, by the function $f(z)$ given by (2.13).

With the aid of Lemma 2, we can similarly prove the following theorem.

THEOREM 4. Let the function $f(z)$ defined by (1.11) be in the class $\mathcal{C}[\alpha, \beta]$ with $0 \leq \alpha \leq 1/2$ and $0 \leq \beta < 1$. Then

$$|D_z^\lambda f(z)| \geq \frac{|z|^{1-\lambda}}{\Gamma(2-\lambda)} \left\{ 1 - \frac{1-\beta}{2(2-\beta)(1-\alpha)(2-\lambda)} |z| \right\}$$

and

$$|D_z^\lambda f(z)| \leq \frac{|z|^{1-\lambda}}{\Gamma(2-\lambda)} \left\{ 1 + \frac{1-\beta}{2(2-\beta)(1-\alpha)(2-\lambda)} |z| \right\}$$

for $0 \leq \lambda < 1$ and $z \in \mathcal{U}$. The results are sharp for the function $f(z)$ defined by

$$D_z^\lambda f(z) = \frac{z^{1-\lambda}}{\Gamma(2-\lambda)} \left\{ 1 - \frac{1-\beta}{2(2-\beta)(1-\alpha)(2-\lambda)} z \right\}$$

or, equivalently, for the function $f(z)$ given by (2.17).

3. A GENERALIZED FRACTIONAL INTEGRAL OPERATOR

We now recall the following definition of a generalized fractional integral operator introduced by Srivastava et al. [14] (see also [13, p. 344]).

DEFINITION 4. For real numbers $\eta > 0, \gamma$, and δ , the generalized fractional integral operator $I_{0,z}^{\eta,\gamma,\delta}$ of order η is defined, for a function $f(z)$, by

$$I_{0,z}^{\eta,\gamma,\delta} f(z) = \frac{z^{-\eta-\gamma}}{\Gamma(\eta)} \int_0^z (z-\zeta)^{\eta-1} F\left(\eta+\gamma, -\delta; \eta; 1-\frac{\zeta}{z}\right) f(\zeta) d\zeta,$$

where $f(z)$ is an analytic function in a simply-connected region of the z -plane containing the origin with the order

$$f(z) = O(|z|^\epsilon), \quad (z \rightarrow 0)$$

$$\epsilon > m \quad (0, \gamma - \delta) - 1$$

and

$$F(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!}, \quad (z \in \mathcal{U}),$$

$(v)_n$ being the Pochhammer symbol defined by

$$(v)_n = \frac{\Gamma(v+n)}{\Gamma(v)} = \begin{cases} 1, & (n=0) \\ v(v+1) \cdots (v+n-1), & (n \in \mathbb{N}) \end{cases}$$

provided further that the multiplicity of $(z-\zeta)^{\eta-1}$ is removed by requiring $\log(z-\zeta)$ to be real when $z-\zeta > 0$.

ReMarK 1. For $\gamma = -\eta$, we readily have

$$I_{0,z}^{\eta,-\eta,\delta} f(z) = D_z^{-\eta} f(z), \quad (\eta > 0),$$

where the fractional integral operator $D_z^{-\lambda} (\lambda > 0)$ is given by Definition 1 .

In order to prove our results involving the generalized fractional integral operator $I_{0,z}^{\eta,\gamma,\delta}$, we shall require the following lemma due to Srivastava et al. [14].

LEMMA 3. If $\eta > 0$ and $\kappa > \gamma - \delta - 1$, then

$$I_{0,z}^{\eta,\gamma,\delta} z^\kappa = \frac{\Gamma(\kappa+1)\Gamma(\kappa-\gamma+\delta+1)}{\Gamma(\kappa-\gamma+1)\Gamma(\kappa+\eta+\delta+1)} z^{\kappa-\gamma}$$

With the aid of Lemma 3, we prove the following theorem.

THEOREM 5. Let $\eta > 0, \gamma < 2, \eta + \delta > -2, \gamma - \delta < 2$, and $\gamma(\eta + \delta) \leq 3\eta$. If the function $f(z)$ defined by (1.11) is in the class $\mathcal{R}[\alpha, \beta]$ with $0 \leq \alpha \leq 1/2$ and $0 \leq \beta < 1$, then

$$\left| I_{0,z}^{\eta,\gamma,\delta} f(z) \right| \geq \frac{\Gamma(2-\gamma+\delta)|z|^{1-\gamma}}{\Gamma(2-\gamma)\Gamma(2+\eta+\delta)} \left\{ 1 - \frac{(1-\beta)(2-\gamma+\delta)}{(2-\beta)(1-\alpha)(2-\gamma)(2+\eta+\delta)} |z| \right\}$$

and

$$\left| I_{0,z}^{\eta,\gamma,\delta} f(z) \right| \leq \frac{\Gamma(2-\gamma+\delta)|z|^{1-\gamma}}{\Gamma(2-\gamma)\Gamma(2+\eta+\delta)} \left\{ 1 + \frac{(1-\beta)(2-\gamma+\delta)}{(2-\beta)(1-\alpha)(2-\gamma)(2+\eta+\delta)} |z| \right\}$$

for $z \in \mathcal{U}_0$, where

$$\mathcal{U}_0 = \begin{cases} \mathcal{U}, & (\gamma \leq 1) \\ \mathcal{U} \setminus \{0\}, & (\gamma > 1) \end{cases}$$

The equalities in (3.5) and (3.6) are attained by the function $f(z)$ given by (2.13).

Proof. By using Lemma 3, we have

$$I_{0,z}^{\eta,\gamma,\delta} f(z) = \frac{\Gamma(2-\gamma+\delta)}{\Gamma(2-\gamma)\Gamma(2+\eta+\delta)} z^{1-\gamma} - \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(n-\gamma+\delta+1)}{\Gamma(n-\gamma+1)\Gamma(n+\eta+\delta+1)} a_n z^{n-\gamma}.$$

Letting

$$\begin{aligned} H(z) &= \frac{\Gamma(2-\gamma)\Gamma(2+\eta+\delta)}{\Gamma(2-\gamma+\delta)} z^\gamma I_{0,z}^{\eta,\gamma,\delta} f(z) \\ &= z - \sum_{n=2}^{\infty} h(n) a_n z^n \end{aligned}$$

where

$$h(n) = \frac{(2-\gamma+\delta)_{n-1} n!}{(2-\gamma)_{n-1} (2+\eta+\delta)_{n-1}}, \quad (n \in \mathbb{N} \setminus \{1\}),$$

we can see that $h(n)$ is nonincreasing for integers $n(n \geq 2)$, and we have

$$0 < h(n) \leq h(2) = \frac{2(2-\gamma+\delta)}{(2-\gamma)(2+\eta+\delta)}.$$

Therefore, by using (2.9) and (3.11), we have

$$\begin{aligned} |H(z)| &\geq |z| - h(2)|z|^2 \sum_{n=2}^{\infty} a_n \\ &\geq |z| - \frac{(1-\beta)(2-\gamma+\delta)}{(2-\beta)(1-\alpha)(2-\gamma)(2+\eta+\delta)} |z|^2 \end{aligned}$$

and

$$\begin{aligned} |H(z)| &\leq |z| + h(2)|z|^2 \sum_{n=2}^{\infty} a_n \\ &\leq |z| + \frac{(1-\beta)(2-\gamma+\delta)}{(2-\beta)(1-\alpha)(2-\gamma)(2+\eta+\delta)} |z|^2. \end{aligned}$$

This completes the proof of Theorem 5 .

Similarly, by applying Lemma 2 (instead of Lemma 1) to the function $f(z)$ belonging to the class $\mathcal{C}[\alpha, \beta]$, we can derive the following theorem.

THEOREM 6. Let $\eta > 0, \gamma < 2, \eta + \delta > -2, \gamma - \delta < 2$, and $\gamma(\eta + \delta) \leq 3\eta$. If the function $f(z)$ defined by (1.11) is in the class $\mathcal{C}[\alpha, \beta]$ with $0 \leq \alpha \leq 1/2$ and $0 \leq \beta < 1$, then

$$\left| I_{0,z}^{\eta,\gamma,\delta} f(z) \right| \geq \frac{\Gamma(2-\gamma+\delta)|z|^{1-\gamma}}{\Gamma(2-\gamma)\Gamma(2+\eta+\delta)} \left\{ 1 - \frac{(1-\beta)(2-\gamma+\delta)}{2(2-\beta)(1-\alpha)(2-\gamma)(2+\eta+\delta)} |z| \right\}$$

and

$$\left| I_{0,z}^{\eta,\gamma,\delta} f(z) \right| \leq \frac{\Gamma(2-\gamma+\delta)|z|^{1-\gamma}}{\Gamma(2-\gamma)\Gamma(2+\eta+\delta)} \left\{ 1 + \frac{(1-\beta)(2-\gamma+\delta)}{2(2-\beta)(1-\alpha)(2-\gamma)(2+\eta+\delta)} |z| \right\}$$

for $z \in \mathcal{U}_0$, where \mathcal{U}_0 is defined by (3.7). The equalities in (3.14) and (3.15) are attained by the function $f(z)$ given by (2.17).

REMARK 2 . For $\beta = \alpha$, Theorem 1 and Theorem 3 would reduce immediately to the corresponding results obtained by Owa and Al-Bassam [15].

ReMARK 3. In view of Remark 1, upon setting $\gamma = -\eta = -\lambda$ in Theorems 5 and 6 , we get the assertions of Theorems 1 and 2 , respectively.

4. REFERENCES

1. M. Abramowitz and I. A. Stegun (Eds.), Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, Dover Publications Inc., New York, 1965.
2. A. A. Attiya, Some applications of Mittag-Leffler function in the unit disk, Filomat 30(7) (2016), 2075-2081. <https://doi.org/10.2298/FIL1607075A>
3. D. Bansal and J. K. Prajapat, Certain geometric properties of the Mittag-Leffler functions, Complex Var. Elliptic Equ. 61(3) (2016), 338-350. <https://doi.org/10.1080/17476933.2015.1079628>
4. T. Bulboacă and G. Murugusundaramoorthy, Univalent functions with positive coefficients involving Pascal distribution series, Commun. Korean Math. Soc. 35(3) (2020), 867 – 877. <https://doi.org/10.4134/CKMS.c190413>
5. N. E. Cho, S. Y. Woo and S. Owa, Uniform convexity properties for hypergeometric functions, Fract. Calc. Appl. Anal. 5(3) (2002), 303-313.
6. P. L. Duren, Univalent Functions, Grundlehren der Mathematischen Wissenschaften Series 259 , Springer Verlag, New York, 1983. [7] M. El-Deeb, T. Bulboacă and J. Dziok, Pascal distribution series connected with certain subclasses of univalent functions, Kyungpook Math. J. **59** (2019), 301-314. <https://doi.org/10.5666/KMJ.2019.59.2.301>

7. B. A. Frasin, An application of an operator associated with generalized Mittag-Leffler function, Konuralp J. Math. 7(1) (2019), 199-202.
8. B. A. Frasin, T. Al-Hawary and F. Yousef, Some properties of a linear operator involving generalized Mittag-Leffler function, Stud. Univ. Babeş-Bolyai Math. 65(1) (2020), 67 – 75. <https://doi.org/10.24193/subbmath.2020.1.06>
9. B. A. Frasin, T. Al-Hawary and F. Yousef, Necessary and sufficient conditions for hypergeometric functions to be in a subclass of analytic functions, Afr. Mat. 30(1-2) (2019), 223-230. <https://doi.org/10.1007/s13370-018-0638-5>
10. H. J. Haubold, A. M. Mathai and R. K. Saxena, Mittag-Leffler functions and their applications, J. Appl. Math. 2011(2011), Article ID 298628.
<https://doi.org/10.1155/2011/298628>
11. V. Kiryakova, Generalized Fractional Calculus and Applications, Pitman Research Notes in Mathematics Series 301, Longman Scientific & Technical, Harlow, John Wiley & Sons, Inc., New York, 1994.
12. E. Merkes and B. T. Scott, Starlike hypergeometric functions, Proc. Amer. Math. Soc. 12 (1961), 885-888.
13. G. M. Mittag-Leffler, Sur la nouvelle fonction $E(x)$, C. R. Acad. Sci. Paris 137(1903), 554 – 558.
14. G. Murugusundaramoorthy, Subordination results for spiral-like functions associated with the Srivastava-Attiya operator, Integral Transforms Spec. Funct. 23(2) (2012), 97-103. <https://doi.org/10.1080/10652469.2011.562501>
15. G. Murugusundaramoorthy, D. Răducanu and K. Vijaya, A class of spirallike functions defined by Ruscheweyh-type q -difference operator, Novi Sad J. Math. 49(2)(2019), 59 – 71. <https://doi.org/10.30755/NSJOM.08284>
16. G. Murugusundaramoorthy, K. Vijaya and S. Porwal, Some inclusion results of certain subclass of analytic functions associated with Poisson distribution series, Hacet. J. Math. Stat. 45(4) (2016), 1101-1107. <https://doi.org/10.15672/HJMS.20164513110>
17. G. Murugusundaramoorthy, Subclasses of starlike and convex functions involving Poisson distribution series, Afr. Mat. 28(2017), 1357-1366.
<https://doi.org/10.1007/s13370-017-0520-x>

18. S. Porwal and M. Kumar, A unified study on starlike and convex functions associated with Poisson distribution series, Afr. Mat. 27(5) (2016), 1021-1027.
<https://doi.org/10.1007/s13370-016-0398-z>
19. M. S. Robertson, On the theory of univalent functions, Ann. of Math. (2) 37(2) (1936), 374-408.
20. H. Silverman, Starlike and convexity properties for hypergeometric functions, J. Math. Anal. Appl. 172 (1993), 574-581. <https://doi.org/10.1006/jmaa.1993.1044>
21. H. M. Srivastava, G. Murugusundaramoorthy and S. Sivasubramanian, Hypergeometric functions in the parabolic starlike and uniformly convex domains, Integral Transforms Spec. Funct. 18 (2007), 511 – 520. <https://doi.org/10.1080/10652460701391324>
22. L. Spáček, Contribution à la théorie des fonctions univalentes, Časopis Pro Pěstování Matematiky 62(1932), 12 – 19.
23. A. Swaminathan, Certain sufficient conditions on Gaussian hypergeometric functions, Journal of Inequalities in Pure and Applied Mathematics 5(4) (2004), Article ID 83, 10 pages.
24. A. Wiman, Über die nullstellen der funktionen $E_{\alpha}(x)$, Acta Math. 29 (1905), 217-134.
25. T. R. Prabhakar, A single integral equation with a generalized Mittag – Leffler function in the kernel, Yokohama Math. J. 19 (1971), 7 – 15.
26. Salah, J. and Darus, M., A note on Generalized Mittag-Leffler function and Application, Far East Journal of Mathematical Sciences (FJMS) 48 (1), 33–46, 2011.
27. [Omar Jawabreh](#), [Ahmad Abdel Qader](#), [Jamal Salah](#), [Khaled Al Mashrafi](#), [Emad Al Dein](#) [AL Fahmawee](#), [Basel J. A. Ali](#), Fractional Calculus Analysis of Tourism Mathematical Model, Progr. Fract. Differ. Appl. Vol. 9, No. S1, pp. 1 -11 202) DOI: 10.18576/pfda/09S101 URL: <https://www.naturalspublishing.com/Article.asp?ArtcID=26373>
28. Jamal Salah, Maryam Al Hashmi and Khaled Al Mashrafi, Some propositions on mathematical models of population growth, International Journal of Latest Trends in Engineering and Technology (IJLTET)Special Issue - ICTIMESH - 2022, 64-69.
URL: https://www.ijltet.org/journal_details.php?id=978&j_id=5050
29. Jamal Salah, Hameed Ur Rehman, and, Iman Al- Buwaiqi, Inclusion Results of a Generalized Mittag-Leffler-Type Poisson Distribution in the k-Uniformly Janowski

- Starlike and the k -Janowski Convex Functions, *Mathematics and Statistics*, Vol. 11, No. 1, pp. 22-27, 2023.
DOI: 10.13189/ms.2023.110103
30. Jamal Salah, Riemann and Big Bang Hypotheses, *NEUROQUANTOLOGY*, Vol. 20, No. 16, pp. 1592-1597, 2022. DOI: 10.48047/NQ.2022.20.16. NQ880156
 31. Jamal Salah, Hameed Ur Rehman, and, Iman Al- Buwaiqi, Subclasses of Spiral-Like Functions Associated with the Generalized Mittag-Leffler Function, *NEUROQUANTOLOGY*, Vol. 20, No. 15, pp. 4784-4799, 2022. DOI: 10.14704/NQ.2022.20.15. NQ88484
 32. Jamal Salah, Maryam Al Hashmi, Hameed Ur Rehman, Khaled Al Mashrafi, Modified Mathematical Models in Biology by the Means of Caputo Derivative of A Function with Respect to Another Exponential Function, *Mathematics and Statistics*, Vol. 10, No. 6, pp. 1194 - 1205, 2022. DOI: 10.13189/ms.2022.100605
 33. Jamal Salah, Hameed Ur Rehman, and, Iman Al- Buwaiqi, The Non-Trivial Zeros of the Riemann Zeta Function through Taylor Series Expansion and Incomplete Gamma Function, *Mathematics and Statistics*, vol. 10, no. 2, pp. 410-418, 2022, DOI: 10.13189/ms.2022.100216.
 34. Jamal Salah, Some Remarks and Propositions on Riemann Hypothesis, *Mathematics and Statistics*, vol. 9, no. 2, pp. 159-165, 2021, DOI: 10.13189/ms.2021.090210.
 35. Hameed Ur Rehman, Malina Darus, and Jamal Salah, Generalizing Certain Analytic Functions Correlative to the n -th Coefficient of Certain Class of Bi-Univalent Functions, *Hindawi Journal of Mathematics* Volume 2021, Article ID 6621315, 14 pages DOI: [10.1155/2021/6621315](https://doi.org/10.1155/2021/6621315)
 36. Salah, J., (2021). Applications of differential Subordination and Superordination Involving Certain Fractional Operator. *Current Topics on Mathematics and Computer Science*, vol. 1, pp. 7–16, 2021 DOI: [10.9734/bpi/ctmcs/v1/9050D](https://doi.org/10.9734/bpi/ctmcs/v1/9050D)
 37. Jamal Y. Mohammad Salah, The Consequence of the Analytic Continuity of Zeta Function Subject to an Additional Term and a Justification of the Location of the Non-Trivial Zeros, *International Journal of Science and Research (IJSR)*, Vol. 9, Issue 3, March 2020, pp. 1566-1569, March 2020, URL: https://www.ijsr.net/get_abstract.php?paper_id=SR20328201050
 38. Jamal Y. Mohammad Salah, An Alternative perspective to Riemann Hypothesis, *PSYCHOLOGY AND EDUCATION*, vol. 57, no. 9, pp. 1278-1281, 2020, URL: [file:///C:/Users/drjamals/Downloads/454-Article%20Text-738-1-10-20210128%20\(1\).pdf](file:///C:/Users/drjamals/Downloads/454-Article%20Text-738-1-10-20210128%20(1).pdf)
 39. Jamal Y. Mohammad Salah, Two Conditional proofs of Riemann Hypothesis, *International Journal of Sciences: Basic and Applied Research (IJSBAR)*, vol. 49, no. 1, pp. 74-83, 2020, URL: <https://gssrr.org/index.php/JournalOfBasicAndApplied/article/view/10720>
 40. Jamal Salah, TWO NEW EQUIVALENT STATEMENTS TO RIEMANN HYPOTHESIS, *Far East Journal of Mathematical Sciences (FJMS)* Volume 118, Issue1, 2019, Pages 1- 8. Sept 2019. DOI: [10.17654/MS118010001](https://doi.org/10.17654/MS118010001)
 41. Jamal Y. Salah, A NEW SUBCLASS OF UNIVALENT FUNCTIONS DEFINED BY THE MEANS OF JAMAL OPERATOR, *Far East Journal of Mathematical Sciences (FJMS)* Vol. 108, No. 2, 2018, pp. 389-399, 2018, Dec 2018, DOI: [10.17654/MS108020389](https://doi.org/10.17654/MS108020389).
 42. Hameed Ur Rehman, Maslina Darus and **Jamal Salah**, Graphing Examples of Starlike and Convex Functions of order β , *Appl. Math. Inf. Sci.* vol. 12, No. 3, pp. 509-515, 2018,

DOI:10.18576/amis/120305

43. Jamal Salah, Hameed Ur Rehman and Maslina Darus, A Note on Caputo's derivative interpretation in Economy, J. Appl. Math, pp. 1-7, 2018, DOI: 10.1155/2018/1260240.
44. Hameed Ur Rehman, Maslina Darus and Jamal Salah, Normalization of the generalized K– Mittag-Leffler function and ratio to its sequence of partial sums, BSG Proceedings, vol. 25, 2018, pp. 78-94.
45. Hameed Ur Rehman, Maslina Darus and Jamal Salah, COEFFICIENT PROPERTIES INVOLVING THE GENERALIZED K–MITTAG-LEFFLER FUNCTIONS, Trans. J. Math. Mecha. (TJMM) 9 (2017), No. 2, 155-164
46. Jamal Y. Salah. Closed-to-Convex Criterion Associated to the Modified Caputo's fractional Calculus Derivative Operator. Far East Journal of Mathematical Sciences (FJMS). Vol. 101, No. 1, pp. 55-59, 2017, DOI: 10.17654/MS101010055.
47. Jamal Y. Salah. Properties of the Modified Caputo's Derivative Operator for certain analytic functions. International Journal of Pure and Applied Mathematics. September 2016, Vol. 109, No. 3, pp. 665 – 671, 2014, DOI: 10.12732/ijpam. v109i3.14
48. Jamal Salah. A Note on the Modified Caputo's Derivative Operator. Far East Journal of Mathematical Sciences (FJMS). Vol. 100, No. 4, pp. 609-615, 2016,

DOI: 10.17654/MS100040609

49. S. Venkatesh, Jamal Salah, G. Sethuraman. "Some Results on E – Cordial Graphs", *International Journal of Mathematical Trends and Technology (IJMTT)*. V7:121-125 March 2014.
DOI: 10.14445/22315373/IJMTT-V7P516
50. Jamal Y. Salah, A Note on Gamma Function, International Journal of Modern Sciences and Engineering Technology (IJMSET), vol. 2, no. 8, pp. 58-64, 2015,
51. Jamal Y. Salah, On Riemann Hypothesis and Robin's Inequality, International Journal of Scientific and Innovative Mathematical Research (IJSIMR), vol. 3, no. 4, pp. 9-14, April 2015,
URL: <https://www.arcjournals.org/pdfs/ijsimr/v3-i4/4.pdf>
52. Jamal Y. Salah, **A Note on Riemann Zeta Function**, IOSR Journal of Engineering (IOSRJEN), vol. 06, no. 02, pp. 07-16, February 2016,
URL: [http://iosrjen.org/Papers/vol6_issue2%20\(part-3\)/B06230716.pdf](http://iosrjen.org/Papers/vol6_issue2%20(part-3)/B06230716.pdf)
53. Jamal Y. Salah, A NOTE ON THE HURWITZ ZETA FUNCTION, Far East Journal of Mathematical Sciences (FJMS), vol. 101, no. 12, pp. 2677-2683, June 2017,
DOI: [10.17654/MS101122677](https://doi.org/10.17654/MS101122677)
54. Jamal Salah and Venkatesh Srivastava. Inequalities on the Theory of Univalent Function. Journal of Mathematics and System Science, vol. 4, pp. 509-513, 2014.
DOI: 10.17265/2159-5291/2014.07.008
55. Jamal Salah. Neighborhood of a certain family of multivalent functions with negative coefficients. International Journal of Pure and Applied Mathematics. (IJPAM). Vol. 92, No 4, April 2014,
DOI: 10.12732/ijpam. v92i4.14
56. Jamal Salah. Fekete-Szego problems involving certain integral operator. International Journal of Mathematics Trends and Technology. IJMTT, vol. 7, no. 1, pp. 54-60, 2014.
DOI: 10.14445/22315373/IJMTT-V7P508
57. Jamal Salah. Subordination and superordination involving certain fractional operator. Asian Journal of Fuzzy and Applied Mathematics, vol. 1, pp. 98-107, 2013.

URL: <https://www.ajouronline.com/index.php/AJFAM/article/view/724>

58. Jamal Salah. Certain subclass of analytic functions associated with fractional calculus operator. *Trans. J. Math. Mecha*, vol. 3, no. 1, pp. 35-42, 2011.
URL: <http://tjmm.edyopress.ro/journal/11030106.pdf>
59. Jamal Salah. A note on Starlike functions of order α associated with a fractional calculus operator involving Caputo's fractional. *J. Appl. comp Sc. Math*, vol. 5, no. 1, pp. 97- 101, 2011,
URL: https://www.jacsm.ro/view/?pid=10_16
60. Jamal Salah and Maslina Darus. On convexity of general integral operators.
J. Anale. Universitatii. De Vest. Timisoara. Seria Matematica – Informatica XLIX. 1 (2011) 117-124.
61. Jamal Salah and Maslina Darus. A note on generalized Mittag-Leffler and Application. *Far East. Math Sc. (FJMS)*, vol. 48, no. 1, pp. 33-46, 2011.
URL: <http://www.pphmj.com/abstract/5478.htm>
62. Jamal Salah and Maslina Darus. A subclass of uniformly convex functions associated with fractional calculus operator involving Caputo's fractional differentiation. *Acta. Univ. APL*. Vol. 24, pp. 295-306, 2010.
63. D. G. Cantor, "Power series with integral coefficients," *Bulletin of the American Mathematical Society*, vol. 69 , no. 3, pp. 362 – 367, 1963.
64. R. Vein and P. Dale, "Determinants and Their Applications in Mathematical Physics," in *Applied Mathematical Sciences*, vol. 134, Springer, New York, 1999.
65. R. Wilson, "Determinantal criteria for meromorphic functions," *Proceedings of the London Mathematical Society*, vol. s3-4, no. 1, pp. 357-374, 1954.
66. C. Pommerenke, "On the coefficients and Hankel determinants of univalent functions," *Journal of the London Mathematical Society*, vol. s1-41, no. 1, pp. 111-122, 1966. [5] C. Pommerenke, "On the Hankel determinants of univalent functions," *Mathematika*, vol. 14, no. 1, pp. 108-112, 1967.
67. D. Bansal, "Upper bound of second Hankel determinant for a new class of analytic functions," *Applied Mathematics Letters*, vol. 26, no. 1, pp. 103-107, 2013.
68. M. M. Elhosh, "On the second Hankel determinant of univalent functions," *Bull. Malays. Math. Soc.*, vol. 9, no. 1, pp. 23-25, 1986.
69. J. W. Noonan and D. K. Thomas, "On the Hankel determinants of areally mean p-valent functions," *Proceedings of the London Mathematical Society*, vol. s3-25, no. 3, pp. 503-524, 1972.
70. J. W. Noonan, "Coefficient differences and Hankel determinants of areally mean p-valent functions," *Proceedings of American Mathematical Society*, vol. 46, pp. 29-37, 1974.
71. J. W. Noonan and D. K. Thomas, "On the second Hankel determinant of areally mean p-valent functions," *Transactions of the American Mathematical Society*, vol. 223, pp. 337-346, 1976.
72. M. M. Elhosh, "On the second Hankel determinant of close-toconvex functions," *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 9, no. 2, pp. 67-68, 1986.

73. M. Arif, K. I. Noor, and M. Raza, "Hankel determinant problem of a subclass of analytic functions," *Journal of Inequalities and Applications*, vol. 2012, Article ID 22, 2012
74. T. Hayami and S. Owa, "Generalized Hankel determinant for certain classes," *International Journal of Mathematics and Analysis*, vol. 4, no. 49-52, pp. 2573-2585, 2010.
75. T. Hayami and S. Owa, "Applications of Hankel determinant for p-valently starlike and convex functions of order α ," *Far East Journal of Applied Mathematics*, vol. 46, no. 1, pp. 1-23, 2010.
76. T. Hayami and S. Owa, "Hankel determinant for p-valently starlike and convex functions of order α ," *General Math*, vol. 17, no. 4, pp. 29-44, 2009.
77. A. Janteng, S. A. Halim, and M. Darus, "Hankel determinant for starlike and convex functions," *International Journal of Mathematics and Analysis*, vol. 1, no. 13-16, pp. 619-625, 2007.
78. A. K. Mishra and P. Gochhayat, "Second Hankel determinant for a class of analytic functions defined by fractional derivative," *International Journal of Mathematics and Mathematical Sciences*, vol. 2008, Article ID 153280, 10 pages, 2008.
79. N. Mohamed, D. Mohamad, and S. Cik Soh, "Second Hankel determinant for certain generalized classes of analytic functions," *International Journal of Mathematics and Analysis*, vol. 6, no. 17-20, pp. 807-812, 2012.
80. G. Murugusundaramoorthy and N. Magesh, "Coefficient inequalities for certain classes of analytic functions associated with Hankel determinant," *Bulletin of Mathematical Analysis and Applications*, vol. 1, no. 3, pp. 85-89, 2009.
81. K. I. Noor, "Higher order close-to-convex functions," *Japanese Journal of Mathematics*, vol. 37, no. 1, pp. 1-8, 1992.
82. K. I. Noor, "On the Hankel determinant problem for strongly close-to-convex functions," *Journal of Natural Geometry*, vol. 11, no. 1, pp. 29-34, 1997.
83. K. I. Noor, "On certain analytic functions related with strongly close-to-convex functions," *Applied Mathematics and Computation*, vol. 197, no. 1, pp. 149-157, 2008. [23] K. I. Noor and S. A. Al-Bany, "On Bazilevic functions," *International Journal of Mathematics and Mathematical Sciences*, vol. 10, no. 1, 88 pages, 1987.
84. K. I. Noor, "On analytic functions related with functions of bounded boundary rotation," *Commentarii Mathematici Universitatis Sancti Pauli*, vol. 30, no. 2, pp. 113-118, 1981.
85. K. I. Noor, "On meromorphic functions of bounded boundary rotation," *Caribb. J. Math.*, vol. 1, no. 3, pp. 95-103, 1982.
86. K. I. Noor, "Hankel determinant problem for the class of functions with bounded boundary rotation," *Revue Roumaine de Mathématiques Pures et Appliquées*, vol. 28, no. 8, pp. 731-739, 1983.
87. K. I. Noor and I. M. A. Al-Naggar, "On the Hankel determinant problem," *Journal of Natural Geometry*, vol. 14, no. 2, pp. 133-140, 1998.
88. R. M. Ali, S. K. Lee, V. Ravichandran, and S. Supramaniam, "The Fekete-Szegő coefficient functional for transforms of analytic functions," *Bulletin of the Iranian Mathematical Society*, vol. 35, no. 2, pp. 119-142, 2009.
89. R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Coefficient bounds for p₋-valent functions," *Applied Mathematics and Computation*, vol. 187, no. 1, pp. 35-46, 2007. W. C. Ma and D. Minda, "A united treatment of some special classes of univalent functions," in *Proceedings of the Conference on Complex Analysis*, Tianjin, 1992.
90. R. K. Raina and J. Sokół, "On coefficient estimates for a certain class of starlike functions, Hacet," *Journal of Mathematics and Statistics*, vol. 44, no. 6, pp. 1427-1433, 2015.

91. G. Murugusundaramoorthy and T. Bulboacă, "Hankel determinants for new subclasses of analytic functions related to a shell shaped region," *Mathematics*, vol. 8, no. 6, article 1041, 2020.
92. S. Gandhi, "Radius Estimates for Three Leaf Function and Convex Combination of Starlike Functions," in *Mathematical Analysis I: Approximation Theory. ICRAPAM 2018*, vol. 306 of *Springer Proceedings in Mathematics & Statistics*, Springer, Singapore.
93. S. N. Malik, S. Mahmood, M. Raza, S. Farman, and S. Zainab, "Coefficient inequalities of functions associated with petal type domains," *Mathematics*, vol. 6, no. 12, p. 298, 2018.
94. K. Sharma, N. K. Jain, and V. Ravichandran, "Starlike functions associated with a cardioid," *Afrika Matematika*, vol. 27, pp. 923-939, 2016.
95. V. S. Masih and S. Kanas, "Subclasses of starlike and convex functions associated with the Limaçon domain," *Symmetry*, vol. 12, no. 6, p. 942, 2020.
96. Y. Yuzaimi, A. H. Suzeini, and A. B. Akbarally, "Subclass of starlike functions associated with a limaçon," *AIP Conference Proceedings*, vol. 1974, article 030023, 2018.
97. P. L. Duren, *Univalent Functions. Grundlehren der Mathematischen Wissenschaften*, vol. 259, Springer, New York, 1983.
98. U. Grenander and G. Szegő, *Toeplitz Forms and Their Applications. California Monographs in Mathematical Sciences*, University of California Press, Berkeley, 1958.
99. M. Abramowitz and I. A. Stegun (Eds.), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover Publications Inc., New York, 1965.
100. A. A. Attiya, Some applications of Mittag-Leffler function in the unit disk, *Filomat* 30(7) (2016), 2075-2081. <https://doi.org/10.2298/FIL1607075A>
101. D. Bansal and J. K. Prajapat, Certain geometric properties of the Mittag-Leffler functions, *Complex Var. Elliptic Equ.* 61(3) (2016), 338-350. <https://doi.org/10.1080/17476933.2015.1079628>
102. T. Bulboacă and G. Murugusundaramoorthy, Univalent functions with positive coefficients involving Pascal distribution series, *Commun. Korean Math. Soc.* 35(3) (2020), 867 – 877. <https://doi.org/10.4134/CKMS.c190413>
103. N. E. Cho, S. Y. Woo and S. Owa, Uniform convexity properties for hypergeometric functions, *Fract. Calc. Appl. Anal.* 5(3) (2002), 303-313.
104. P. L. Duren, *Univalent Functions, Grundlehren der Mathematischen Wissenschaften Series 259*, Springer Verlag, New York, 1983. [7] M. El-Deeb, T. Bulboacă and J. Dziok, Pascal distribution series connected with certain subclasses of univalent functions, *Kyungpook Math. J.* **59** (2019), 301-314. <https://doi.org/10.5666/KMJ.2019.59.2.301>
105. B. A. Frasin, An application of an operator associated with generalized Mittag-Leffler function, *Konuralp J. Math.* 7(1) (2019), 199-202.
106. B. A. Frasin, T. Al-Hawary and F. Yousef, Some properties of a linear operator involving generalized Mittag-Leffler function, *Stud. Univ. Babeş-Bolyai Math.* 65(1) (2020), 67 – 75. <https://doi.org/10.24193/subbbmath.2020.1.06>
107. B. A. Frasin, T. Al-Hawary and F. Yousef, Necessary and sufficient conditions for hypergeometric functions to be in a subclass of analytic functions, *Afr. Mat.* 30(1-2) (2019), 223-230. <https://doi.org/10.1007/s13370-018-0638-5>
108. H. J. Haubold, A. M. Mathai and R. K. Saxena, Mittag-Leffler functions and their applications, *J. Appl. Math.* 2011(2011), Article ID 298628. <https://doi.org/10.1155/2011/298628>
109. V. Kiryakova, *Generalized Fractional Calculus and Applications*, Pitman Research Notes in Mathematics Series 301, Longman Scientific & Technical, Harlow, John Wiley & Sons, Inc., New York, 1994.

110. E. Merkes and B. T. Scott, Starlike hypergeometric functions, *Proc. Amer. Math. Soc.* 12 (1961), 885-888.
111. G. M. Mittag-Leffler, Sur la nouvelle fonction $E(x)$, *C. R. Acad. Sci. Paris* 137(1903), 554 – 558.
112. G. Murugusundaramoorthy, Subordination results for spiral-like functions associated with the Srivastava-Attiya operator, *Integral Transforms Spec. Funct.* 23(2) (2012), 97-103. <https://doi.org/10.1080/10652469.2011.562501>
113. G. Murugusundaramoorthy, D. Răducanu and K. Vijaya, A class of spirallike functions defined by Ruscheweyh-type q -difference operator, *Novi Sad J. Math.* 49(2)(2019), 59 – 71. <https://doi.org/10.30755/NSJOM.08284>
114. G. Murugusundaramoorthy, K. Vijaya and S. Porwal, Some inclusion results of certain subclass of analytic functions associated with Poisson distribution series, *Hacet. J. Math. Stat.* 45(4) (2016), 1101-1107. <https://doi.org/10.15672/HJMS.20164513110>
115. G. Murugusundaramoorthy, Subclasses of starlike and convex functions involving Poisson distribution series, *Afr. Mat.* 28(2017), 1357-1366. <https://doi.org/10.1007/s13370-017-0520-x>
116. S. Porwal and M. Kumar, A unified study on starlike and convex functions associated with Poisson distribution series, *Afr. Mat.* 27(5) (2016), 1021-1027. <https://doi.org/10.1007/s13370-016-0398-z>
117. M. S. Robertson, On the theory of univalent functions, *Ann. of Math.* (2) 37(2) (1936), 374-408.
118. Salah, Jamal, Hameed Ur Rehman, and Iman Al-Buwaiqi. "The Non-Trivial Zeros of the Riemann Zeta Function through Taylor Series Expansion and Incomplete Gamma Function." *Mathematics and Statistics* 10.2 (2022): 410-418.
119. Rehman, Hameed Ur, Maslina Darus, and Jamal Salah. "Graphing Examples of Starlike and Convex Functions of order β ." *Appl. Math. Inf. Sci.* 12.3 (2018): 509-515.
120. Salah, Jamal Y., and O. M. A. N. Ibra. "Properties of the Modified Caputo's Derivative Operator for certain analytic functions." *International Journal of Pure and Applied Mathematics* 109.3 (2014): 665-671.
121. Jamal Salah and Maslina Darus, On convexity of the general integral operators, *An. Univ. Vest Timis. Ser. Mat. -Inform.* 49(1) (2011), 117-124.
122. Salah, Jamal. "TWO NEW EQUIVALENT STATEMENTS TO RIEMANN HYPOTHESIS." (2019).
123. H. Silverman, Starlike and convexity properties for hypergeometric functions, *J. Math. Anal. Appl.* 172 (1993), 574-581. <https://doi.org/10.1006/jmaa.1993.1044>
124. H. M. Srivastava, G. Murugusundaramoorthy and S. Sivasubramanian, Hypergeometric functions in the parabolic starlike and uniformly convex domains, *Integral Transforms Spec. Funct.* 18 (2007), 511 – 520. <https://doi.org/10.1080/10652460701391324>
125. L. Spaček, Contribution à la théorie des fonctions univalentes, *Časopis Pro Pěstování Matematiky* 62(1932), 12 – 19.
126. Salah, Jamal Y. "A new subclass of univalent functions defined by the means of Jamal operator." *Far East Journal of Mathematical Sciences (FJMS)* Vol 108 (2018): 389-399.
127. Jamal Y. Salah On Riemann Hypothesis and Robin's Inequality. *International Journal of Scientific and Innovative Mathematical Research (IJSIMR)*. Volume (3) 4 (2015) 9-14.

128. Salah, Jamal. "Neighborhood of a certain family of multivalent functions with negative coefficients." *Int. J. Pure Appl. Math* 92.4 (2014): 591-597.
129. Salah, Jamal, and Maslina Darus. "A note on Starlike functions of order α associated with a fractional calculus operator involving Caputo's fractional." *J. Appl. comp Sc. Math* 5.1 (2011): 97-101.
130. J. Salah, Certain subclass of analytic functions associated with fractional calculus operator, *Trans. J. Math. Mech.*, 3 (2011), 35–42. Available from: <http://tjmm.edyopress.ro/journal/11030106.pdf>.
131. Salah, Jamal. "Some Remarks and Propositions on Riemann Hypothesis." *Mathematics and Statistics* 9.2 (2021): 159-165.
132. Salah, Jamal Y. Mohammad. "The consequence of the analytic continuity of Zeta function subject to an additional term and a justification of the location of the non-trivial zeros." *International Journal of Science and Research (IJSR)* 9.3 (2020): 1566-1569.
- 133.
134. A. Swaminathan, Certain sufficient conditions on Gaussian hypergeometric functions, *Journal of Inequalities in Pure and Applied Mathematics* 5(4) (2004), Article ID 83, 10 pages.
135. A. Wiman, Über die nullstellen der funktionen $E_\alpha(x)$, *Acta Math.* 29 (1905), 217-134.
136. Salah, Jamal Y. Mohammad. "An Alternative perspective to Riemann Hypothesis." *PSYCHOLOGY AND EDUCATION* 57.9 (2020): 1278-1281.
137. Salah, Jamal Y. "A note on the Hurwitz zeta function." *Far East Journal of Mathematical Sciences (FJMS)* 101.12 (2017): 2677-2683.
138. Jamal Y. Salah. Closed-to-Convex Criterion Associated to the Modified Caputo's fractional Calculus Derivative Operator. *Far East Journal of Mathematical Sciences (FJMS)*. Vol. 101, No. 1, pp. 55-59, 2017, DOI: 10.17654/MS101010055.
139. Jamal Y. Salah, A Note on Riemann Zeta Function, *IOSR Journal of Engineering (IOSRJEN)*, vol. 06, no. 02, pp. 07-16, February 2016, URL: [http://iosrjen.org/Papers/vol6_issue2%20\(part-3\)/B06230716.pdf](http://iosrjen.org/Papers/vol6_issue2%20(part-3)/B06230716.pdf)
140. Jamal Y. Salah, A Note on Gamma Function, *International Journal of Modern Sciences and Engineering Technology (IJMSET)*, vol. 2, no. 8, pp. 58-64, 2015.
141. Salah, Jamal, and S. Venkatesh. "Inequalities on the Theory of Univalent Functions." *Journal of Mathematics and System Science* 4.7 (2014).
142. T. R. Prabhakar, A single integral equation with a generalized Mittag – Leffler function in the kernel, *Yokohama Math. J.* 19 (1971), 7 – 15.
143. [Omar Jawabreh](#), [Ahmad Abdel Qader](#), [Jamal Salah](#), [Khaled Al Mashrafi](#), [Emad Al Dein AL Fahmawee](#), [Basel J. A. Ali](#), Fractional Calculus Analysis of Tourism Mathematical Model, *Progr. Fract. Differ. Appl.* Vol. 9, No. S1, pp. 1 -11 202) DOI: 10.18576/pfda/09S101 URL: <https://www.ajouronline.com/index.php/AJFAM/article/view/724>
144. Jamal Salah. Subordination and superordination involving certain fractional operator. *Asian Journal of Fuzzy and Applied Mathematics*, vol. 1, pp. 98-107, 2013. URL: <https://www.ajouronline.com/index.php/AJFAM/article/view/724>

145. Salah, Jamal Y. Mohammad. "Two Conditional proofs of Riemann Hypothesis." *International Journal of Sciences: Basic and Applied Research (IJSBAR)* 49.1 (2020): 74-83.
146. Salah, Jamal. "Fekete-Szego problems involving certain integral operator." *International Journal of Mathematics Trends and Technology-IJMTT* 7 (2014).